


MTM5135-Nonlinear Dynamical Systems

Gökhan Göksu, PhD


Week 11



Comparison Functions




$$\begin{cases} \alpha \text{ continuous} \\ \alpha(0) = 0 \\ \alpha(s) > 0, \forall s > 0 \end{cases}$$



$$\begin{cases} \alpha \in \mathcal{PD} \\ \alpha \text{ nondecreasing} \end{cases}$$



$$\begin{cases} \alpha \in \mathcal{K} \\ \lim_{s \rightarrow \infty} \alpha(s) = \infty \end{cases}$$



$$\begin{cases} \beta(\cdot, t) \in \mathcal{K}, \forall t \geq 0 \\ \beta(s, \cdot) \text{ nonincreasing}, \forall s \geq 0 \\ \lim_{t \rightarrow \infty} \beta(s, t) = 0, \forall s \geq 0 \end{cases}$$

Comparison Functions: Examples

Example

- ▶ $\alpha(s) = \frac{1}{1+s^c}$, for any $c > 0$
- ▶ $\alpha(s) = \frac{s^c}{1+s^c}$, for any $c > 0$
- ▶ $\alpha(s) = \tan^{-1}(s)$
- ▶ $\alpha(s) = \text{sat}(s) = \begin{cases} s, & \text{if } |s| \leq 1 \\ \text{sgn}(s), & \text{if } |s| > 1 \end{cases}$
- ▶ $\alpha(s) = \text{sat}(s; a, b) = a + \frac{2(b-a)}{\pi} \tan^{-1}(s)$
- ▶ $\alpha(s) = s^c$, for any $c > 0$
- ▶ $\alpha(s) = \min\{s, s^2\}$
- ▶ $\beta(s, r) = \frac{s}{krs+1}$
- ▶ $\beta(s, r) = s^c e^{-r}$

Equivalent Representation of Radial Unboundedness

Lemma (4.3 in [Khalil, 2002])

Let $V : D \rightarrow \mathbb{R}$ be a continuous positive definite function (may not be \mathcal{PD} !) defined on $D \subset \mathbb{R}^n$ that contains the origin. Let $B_r[0] \subset D$ for some $r > 0$. Then, there exist $\alpha_1, \alpha_2 \in \mathcal{K}$ defined on $[0, r]$ such that

$$\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|)$$

for all $x \in B_r[0]$. If $D = \mathbb{R}^n$, the functions α_1 and α_2 will be defined on $[0, \infty)$ and the foregoing inequality will hold for all $x \in \mathbb{R}^n$. Moreover, if $V(x)$ is radially unbounded, then $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$.

Example

- ▶ $V(x) = x^\top P x \implies \lambda_{\min}(P)|x|^2 \leq x^\top P x \leq \lambda_{\max}(P)|x|^2$
- ▶ Given $x \in \mathbb{R}^n$, $|x|$ denotes its Euclidean norm.

Equivalent Representation of GAS

For a system with no inputs $\dot{x} = f(x)$, there is a well-known notion of global asymptotic stability (for short from now on, GAS, or “0-GAS” when referring to the system with no-inputs $\dot{x} = f(x, 0)$ associated to a given system with inputs $\dot{x} = f(x, u)$ due to Lyapunov, and usually defined in “ ϵ - δ ” terms. It is easy to show that the standard definition of GAS (0-GAS) in previous lecture is in fact equivalent to the existence of $\beta \in \mathcal{KL}$ satisfying the following, along the solutions of $\dot{x} = f(x)$ ($\dot{x} = f(x, 0)$)

$$|x(t, x_0)| \leq \beta(|x_0|, t), \quad \forall x_0 \in \mathbb{R}^n, \forall t \geq 0.$$

Observe that, since β decreases on t , we have, in particular:

$$|x(t, x_0)| \leq \beta(|x_0|, 0), \quad \forall x_0 \in \mathbb{R}^n, \forall t \geq 0.$$

which provides the stability part of the GAS definition while

$$|x(t, x_0)| \leq \beta(|x_0|, t) \xrightarrow[t \rightarrow \infty]{} 0, \quad \forall x_0 \in \mathbb{R}^n,$$

which is the attractivity (convergence to the equilibrium point) part of the GAS definition.

Note: From now on, unless written explicitly, the solutions $x(t, x_0)$ or $x(t, x_0, u)$ for $\dot{x} = f(x)$ and $\dot{x} = f(x, u)$, respectively, will be written in short as $x(t)$ to avoid cumbersome notation!

Equivalent Lyapunov Theorem for Stability

Theorem (4.8 in [Khalil, 2002])

Let $x = 0$ be an equilibrium point for $\dot{x} = f(x)$ and $D \subset \mathbb{R}^n$ be a domain containing the origin. Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable function such that

$$W_1(x) \leq V(x) \leq W_2(x)$$
$$\frac{\partial V}{\partial x} f(x) \leq 0$$

for all $t \geq 0$ and $x \in D$, where W_1 and W_2 are continuous positive definite functions on D . Then, $x = 0$ is stable.

Equivalent Lyapunov Theorem for AS

Theorem (4.9 in [Khalil, 2002])

Let $x = 0$ be an equilibrium point for $\dot{x} = f(x)$ and $D \subset \mathbb{R}^n$ be a domain containing the origin. Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable function such that

$$W_1(x) \leq V(x) \leq W_2(x)$$

$$\frac{\partial V}{\partial x} f(x) \leq -W_3(x)$$

for all $t \geq 0$ and $x \in D$, where W_1 , W_2 and W_3 are continuous positive definite functions on D . Then, $x = 0$ is **asymptotically** stable. Moreover, if r and c are chosen such that $B_r[0] = \{x \in D \mid |x| \leq r\}$ and $c < \min_{|x|=r} W_1(x)$, then every trajectory starting in $\{x \in B_r[0] \mid W_2(x) \leq c\}$ satisfies

$$|x(t)| \leq \beta(|x(0)|, t), \quad \forall t \geq 0$$

for some $\beta \in \mathcal{KL}$.

Equivalent Lyapunov Theorem for AS




Figure: Geometric representation of sets in Theorem 4.9.

Equivalent Lyapunov Theorem for GAS

Theorem (4.9 in [Khalil, 2002])

Let $x = 0$ be an equilibrium point for $\dot{x} = f(x)$. Let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ ($D = \mathbb{R}^n!$) be a continuously differentiable function such that

$$\alpha_1(|x|) \leq V(x) \leq \alpha_2(|x|)$$
$$\frac{\partial V}{\partial x} f(x) \leq -W_3(x)$$

for all $t \geq 0$ and $x \in \mathbb{R}^n$, where $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ and W_3 is continuous positive definite function on \mathbb{R}^n . Then, $x = 0$ is **GAS**.

Let us remember the Lyapunov theorem for ES/GES shown last week:

Theorem (4.10 in [Khalil, 2002])

Let $x = 0$ be an equilibrium point for $\dot{x} = f(x)$. Let $V : D \rightarrow \mathbb{R}$ be a continuously differentiable function such that

$$k_1|x|^a \leq V(x) \leq k_2|x|^a$$
$$\frac{\partial V}{\partial x} f(x) \leq -k_3|x|^a$$

for all $t \geq 0$ and $x \in D$, where k_1, k_2, k_3 and a are positive constants. Then, $x = 0$ is **ES**. If $D = \mathbb{R}^n$, then $x = 0$ is **GES**.

Nonlinear Systems: 0-GAS \Rightarrow Good Behavior wrt Inputs

- ▶ Car trailer system

- ▶ Video:

- <https://www.youtube.com/watch?v=4jK9H5AB4lM>

- ▶ Aircraft

- ▶ Video:

- <https://www.youtube.com/watch?v=4UfmsqtTGa0>

- ▶ Car active suspension system

- ▶ Video:

- <https://www.youtube.com/watch?v=kRt7H0k8A4k>

- ▶ Building

- ▶ Video:

- <https://www.youtube.com/shorts/rJ72LruGgyU>

Nonlinear Systems: 0-GAS \nRightarrow Good Behavior wrt Inputs

For linear systems $\dot{x} = Ax + Bu$:

- ▶ $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$
- ▶ If A is a Hurwitz matrix ($\text{Re}(\lambda_i(A)) < 0$ for all $i = 1, \dots, n$), then the linear system is 0-GAS.
- ▶ Such a 0-GAS linear system automatically satisfies all reasonable “input-to-state stability” properties [Sontag, 1990]¹:
 - ▶ Bounded inputs \Rightarrow bounded state (BIBS) trajectories
 - ▶ Converging inputs \Rightarrow converging state (CICS) trajectories

This is generally not the case for nonlinear systems $\dot{x} = f(x, u)$!

Example

Consider the scalar system ($n = 1$) with a single input ($m = 1$)

$$\dot{x} = -x + (x^2 + 1)u:$$

- ▶ The system is clearly 0-GAS, since it reduces to $\dot{x} = -x$ when $u \equiv 0$.
- ▶ However, for $u = (2t + 2)^{-1/2}$ and $x_0 = \sqrt{2}$, the system produces unbounded and even diverging state trajectory $x(t) = (2t + 2)^{1/2}$!

¹Mathematical Control Theory: Deterministic Finite Dimensional Systems

Nonlinear Systems: 0-GAS \nRightarrow Good Behavior wrt Inputs




Figure: Diverging state for converging input.

Estimates (Gains) for Linear/Nonlinear Systems

Recall the solution of the linear system $\dot{x} = Ax + Bu$ can be written as:

$$x(t) = e^{At}x(0) + \int_0^t e^{A(t-\tau)}Bu(\tau)d\tau$$

If A is Hurwitz, there exists some $k, \lambda > 0$ such that $\|e^{At}\| \leq ke^{-\lambda t}$ which, in turn, gives the following state estimate

$$\begin{aligned}|x(t)| &\leq k|x(0)|e^{-\lambda t} + \int_0^t ke^{-\lambda(t-\tau)}\|B\||u(\tau)|d\tau \\&\leq k|x(0)|e^{-\lambda t} + k\|B\| \sup_{\tau \in [0,t]} |u(\tau)| e^{-\lambda t} \int_0^t e^{\lambda\tau} d\tau \\&= k|x(0)|e^{-\lambda t} + k\|B\| \sup_{\tau \in [0,t]} |u(\tau)| e^{-\lambda t} \left(\frac{e^{\lambda t} - 1}{\lambda} \right) \\&\leq k|x(0)|e^{-\lambda t} + \frac{k\|B\|}{\lambda} \sup_{\tau \in [0,t]} |u(\tau)| (1 - e^{-\lambda t}) \\&\leq \bar{k}|x(0)|e^{-\lambda t} + \bar{k} \sup_{\tau \in [0,t]} |u(\tau)|\end{aligned}$$

where $\bar{k} = k \cdot \max\{1, \frac{\|B\|}{\lambda}\}$

Estimates (Gains) for Linear/Nonlinear Systems

Motivated with this estimation, for linear systems, three most typical ways of defining “input-to-state stability” in terms of operators $\{L^2, L^\infty\} \rightarrow \{L^2, L^\infty\}$ are as follows:

- ▶ “ $L^\infty \rightarrow L^\infty$ ”: $c|x(t)| \leq |x_0|e^{-\lambda t} + \sup_{\tau \in [0, t]} |u(\tau)|$
- ▶ “ $L^2 \rightarrow L^\infty$ ”: $c|x(t)| \leq |x_0|e^{-\lambda t} + \int_0^t |u(\tau)|^2 d\tau$
- ▶ “ $L^2 \rightarrow L^2$ ”: $c \int_0^t |x(\tau)|^2 d\tau \leq |x_0| + \int_0^t |u(\tau)|^2 d\tau$

The missing case “ $L^\infty \rightarrow L^2$ ” is less interesting, being too restrictive, for practical reasons! Concerning the nonlinear system $\dot{x} = f(x, u)$, in general, under “some” nonlinear coordinate change (see [Sontag, 2004]), we arrive to the following three concepts (or “estimates”) for nonlinear systems:

- ▶ “ $L^\infty \rightarrow L^\infty$ ”: $\alpha(|x(t)|) \leq \beta(|x_0|, t) + \sup_{\tau \in [0, t]} \gamma(|u(\tau)|)$
- ▶ “ $L^2 \rightarrow L^\infty$ ”: $\alpha(|x(t)|) \leq \beta(|x_0|, t) + \int_0^t \gamma(|u(\tau)|) d\tau$
- ▶ “ $L^2 \rightarrow L^2$ ”: $\int_0^t \alpha(|x(\tau)|) d\tau \leq \alpha_0(|x_0|) + \int_0^t \gamma(|u(\tau)|) d\tau$

Here, the functions (which measure the impacts of the state or input) are $\alpha, \alpha_0, \gamma \in \mathcal{K}_\infty$ and $\beta \in \mathcal{KL}$. The “ $L^\infty \rightarrow L^\infty$ ” (and “ $L^2 \rightarrow L^2$ ” as well) estimate leads us to the first concept that of *input-to-state stability (ISS)* whereas “ $L^2 \rightarrow L^\infty$ ” estimate leads us to the second concept that of *integral input-to-state stability (ilSS)*.

ISS and iISS Notions

Definition: Input-to-State Stability (ISS) [Sontag, IEEE TAC, 1989]

The system $\dot{x} = f(x, u)$ is **ISS** if there exist $\beta \in \mathcal{KL}$ and $\nu \in \mathcal{K}_\infty$ such that, for all $x_0 \in \mathbb{R}^n$ and all $u \in \mathcal{U}$,

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \nu(\|u\|), \quad \forall t \geq 0.$$

→ Vanishing transients “proportional” to initial state’s norm

→ Steady-state error “proportional” to input **amplitude**.

Definition: Integral Input-to-State Stability (iISS) [Sontag, SCL, 1998]

The system $\dot{x} = f(x, u)$ is **iISS** if there exist $\beta \in \mathcal{KL}$ and $\nu_1, \nu_2 \in \mathcal{K}_\infty$ such that, for all $x_0 \in \mathbb{R}^n$ and all $u \in \mathcal{U}$,

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \nu_1 \left(\int_0^t \nu_2(|u(s)|) ds \right), \quad \forall t \geq 0.$$

→ Measures the impact of input **energy**.

ISS and iISS Notions: Strengths and Weaknesses

ISS and iISS: Central tools in nonlinear analysis and control:

- ▶ **Theoretical contributions** to: output feedback, optimal control, hybrid systems, predictive control, chaotic systems. . .
- ▶ **Applications** in: robotics, production lines, transportation, bio-chemical networks, control under communication constraints, neuroscience, . . .

ISS	iISS
$\dot{x} = f(x, 0)$ is GAS	$\dot{x} = f(x, 0)$ is GAS
Bounded input \Rightarrow Bounded state	Bounded energy input \Rightarrow Bounded, converging state
Converging input \Rightarrow Converging state	Converging input \nRightarrow Converging state
Cascade: ISS + ISS \Rightarrow ISS	Cascade: iISS + iISS \nRightarrow iISS

In practice, some systems do exhibit robustness for inputs under a certain threshold, but diverge for larger ones.

Strong iISS: halfway between ISS and iISS.


ISS and iISS Notions: Lyapunov Characterizations

- ▶ Part of the success of ISS and iISS is due to their Lyapunov characterizations
- ▶ Lyapunov function candidate (LFC):
 - ▶ $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ continuously differentiable
 - ▶ $V(0) = 0$ and $V(x) > 0$ for all $x \neq 0$
 - ▶ $V(x) \rightarrow \infty$ whenever $|x| \rightarrow \infty$.

Theorem: ISS and iISS Characterization [Sontag, Wang, SCL, 1995] & [Angeli et al., IEEE TAC, 2000]

The system $\dot{x} = f(x, u)$ is ISS (resp. iISS) if and only if there exist a LFC V , $\gamma \in \mathcal{K}_\infty$, and $\alpha \in \mathcal{K}_\infty$ (resp. $\alpha \in \mathcal{PD}$) such that, for all $x \in \mathbb{R}^n$ and all $u \in \mathbb{R}^m$

$$\frac{\partial V}{\partial x} f(x, u) \leq -\alpha(|x|) + \gamma(|u|). \quad (\mathcal{D}\text{-}\mathcal{K}_\infty/\mathcal{PD})$$



ISS Characterization in Implication Form

Theorem: ISS Char. in Implication Form [Sontag, Wang, SCL, 1995]

The system $\dot{x} = f(x, u)$ is ISS if and only if there exist a LFC V , $\chi \in \mathcal{K}$, and $\tilde{\alpha} \in \mathcal{K}$ such that for any $x \in \mathbb{R}^n$ and any $u \in \mathbb{R}^m$

$$|x| \geq \chi(|u|) \implies \frac{\partial V}{\partial x} f(x, u) \leq -\tilde{\alpha}(|x|) \quad (\text{D-IF})$$

Remark

Clearly, $(D-\mathcal{K}_\infty)$ implies $(D\text{-IF})$. Assume now that $(D\text{-IF})$ holds with some $\tilde{\alpha} \in \mathcal{K}$ and $\chi \in \mathcal{K}$. Without loss of generality, one can assume that $\tilde{\alpha} \in \mathcal{K}_\infty$ (see [Lin, Sontag, Wang, SCL, 1995, Remark 4.1]). Let

$$\gamma(r) = \max\{0, \hat{\gamma}(r)\}$$

where

$$\hat{\gamma}(r) = \max \left\{ \frac{\partial V}{\partial x} f(x, u) + \tilde{\alpha}(\chi(|u|)) : |u| \leq r, |x| \leq \chi(r) \right\}.$$

Then $\gamma \in \mathcal{C}$, $\gamma(0) = 0$ and, therefore, $\gamma \in \mathcal{K}_\infty$ by definition. $(D-\mathcal{K}_\infty)$ holds because $\gamma(r) \geq \sup_{|u|=r} \frac{\partial V}{\partial x} f(x, u) + \tilde{\alpha}(\chi(|u|))$ (consider the two separate cases $|x| \geq \chi(|u|)$ and $|x| \leq \chi(|u|)$).

Halfway Between ISS and iISS: Strong iISS Property

Definition: Strong iISS [Chaillet et. al., IEEE TAC, 2014]

The system $\dot{x} = f(x, u)$ is **Strongly iISS** if it is:


- ▶ iISS
- ▶ ISS with respect to small inputs

i.e., if there exist $\beta \in \mathcal{KL}$, $\nu_1, \nu_2, \nu \in \mathcal{K}_\infty$ and **input threshold** $R > 0$ such that, for all $x_0 \in \mathbb{R}^n$ and all $u \in \mathcal{U}$,

$$|x(t; x_0, u)| \leq \beta(|x_0|, t) + \nu_1 \left(\int_0^t \nu_2(|u(s)|) ds \right)$$

$$\|u\| \leq R \Rightarrow |x(t; x_0, u)| \leq \beta(|x_0|, t) + \nu(\|u\|).$$

- ▶ For all $u \in \mathcal{U}$, the solution exists at all times
- ▶ $\int_0^t \nu_2(|u(s)|) ds < \infty \Rightarrow$ bounded and converging state
- ▶ Converging input \Rightarrow converging state
- ▶ $\|u\| \leq R \Rightarrow$ bounded state.



Halfway Between ISS and iISS: Strong iISS Property

Theorem: \mathcal{K} dissipation rate \Rightarrow Strong iISS [Chaillet et. al., IEEE TAC, 2014]

If there exists a LFC $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ satisfying, for all $x \in \mathbb{R}^n$ and all $u \in \mathbb{R}^m$,

$$\frac{\partial V}{\partial x} f(x, u) \leq -\alpha(|x|) + \gamma(|u|).$$

where $\alpha \in \mathcal{K}$ and $\gamma \in \mathcal{K}_{\infty}$, then the system $\dot{x} = f(x, u)$ is **Strongly iISS** with input threshold $R = \gamma^{-1} \circ \alpha(\infty)$.

Equivalently, we can state the following:

Corollary: Non-vanishing dissipation rate \Rightarrow Strong iISS [Chaillet et. al., IEEE TAC, 2014]

If there exists a LFC $V : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ satisfying, for all $x \in \mathbb{R}^n$ and all $u \in \mathbb{R}^m$,

$$\frac{\partial V}{\partial x} f(x, u) \leq -W(x) + \gamma(|u|).$$

where $\gamma \in \mathcal{K}_{\infty}$ and W is continuous positive definite satisfying

$W_{\infty} := \liminf_{|x| \rightarrow \infty} W(x) > 0$, then the system $\dot{x} = f(x, u)$ is **Strongly iISS** with input threshold $R = \gamma^{-1}(W_{\infty})$.

Halfway Between ISS and iISS: Strong iISS Property

However, the converse does not hold:

Counter-example: Strong iISS \nRightarrow \mathcal{K} dissipation rate [Chaillet et. al., IEEE TAC, 2014]

The scalar system

$$\dot{x} = -\frac{x}{1+x^2} \left[1 - |x|(u^2 - |u|) \right],$$


is Strongly iISS. However, for all $\alpha \in \mathcal{K}$ and $\gamma \in \mathcal{K}_\infty$ no differentiable function $V : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ satisfies

$$\frac{\partial V}{\partial x}(x)f(x, u) \leq -\alpha(|x|) + \gamma(|u|).$$

- ▶ iISS: $V_1(x) = \frac{1}{2} \ln(1+x^2)$ gives $\dot{V}_1 \leq -x^2/(1+x^2)^2 + u^2 + |u|$
- ▶ ISS wrt $|u| < 1$: $V_2(x) = x^4/4$ gives $\dot{V}_2 \leq -x^4/(1+x^2)$.

ISS, iISS and Strong iISS Example

Consider the tank with a flat bottom and not necessarily constant cross-section [Daskovski, IFAC POL, 2019].



ISS, iISS and Strong iISS Example

By Toricelli's law, the level x of the liquid changes with time due to the inflow and outflow can be described by the following differential equation:

$$\dot{x} = -\frac{a(x)\mu\sqrt{2gx}}{A(x)} + \frac{u}{A(x)}$$

where

$A : [0, \infty) \rightarrow (0, \infty)$ is the cross-section area of the tank at the height x

$a : [0, \infty) \rightarrow [0, \infty)$ is the area of the hole, that in general may also depend on x

u is the rate of inflow to the tank

Let us consider the following three cases:

- Suppose that $a(x) = a = (\mu\sqrt{2g})^{-1}$ and $A(x) = 1 + \sqrt[4]{x}$. The ISS property can be verified by considering $V(x) = |x| = x$. For this V , we have

$$\dot{V} = -\frac{\sqrt{x}}{1 + \sqrt[4]{x}} + \frac{u}{1 + \sqrt[4]{x}}$$

Now, observe that we have

$$|x| \geq 4u^2 \implies \dot{V} \leq -\frac{\sqrt{x}}{2(1 + \sqrt[4]{x})}$$

and since the function $\tilde{\alpha}(s) = \frac{\sqrt{s}}{2(1 + \sqrt[4]{s})}$ and $\chi(s) = 4s^2$ are two class \mathcal{K}_∞ functions, we can conclude that the system is ISS.

ISS, iISS and Strong iISS Example

Let us consider the following three cases (continued):


- Suppose that $a(x) = \frac{1}{1+x}$ and $A(x) = A = \mu\sqrt{2g}$. The iISS property can be verified by considering $V(x) = |x| = x$. For this V , we have

$$\dot{V} \leq -\frac{\sqrt{x}}{1+x} + \frac{u}{A}$$

Since the function $\alpha(s) = \frac{\sqrt{s}}{1+s}$ is a class \mathcal{PD} function whereas $\gamma(s) = \frac{s}{A}$ is a class \mathcal{K}_∞ function, we can conclude that the system is iISS.

- (Exercise) Show that the system is Strong iISS when $a(x) = \frac{1}{\sqrt{1+x}}$ and $A(x) = A = \mu\sqrt{2g}$.

Cascades of ISS or iISS Systems



- ▶ ISS is naturally preserved in cascade [Sontag, EJC, 1995]
- ▶ iISS is **not** preserved by cascade [Panteley, Loria, Automatica, 2001] & [Arcak et al., SICON, 2002].

Theorem [Chaillet, Angeli, SCL, 2008]


Let V_1 and V_2 be two Lyapunov functional candidates. Assume that there exist $\gamma_1, \gamma_2 \in \mathcal{K}$, and $\alpha_1, \alpha_2 \in \mathcal{PD}$ such that, for all $(x_1, x_2) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$ and all $u \in \mathbb{R}^m$,

$$\begin{aligned}\frac{\partial V_1}{\partial x_1} f_1(x_1, x_2) &\leq -\alpha_1(|x_1|) + \gamma_1(|x_2|) \\ \frac{\partial V_2}{\partial x_2} f_2(x_2, u) &\leq -\alpha_2(|x_2|) + \gamma_2(|u|).\end{aligned}$$

If $\gamma_1(s) = \mathcal{O}_{s \rightarrow 0^+}(\alpha_2(s))$, then the cascade is **iISS**.

- ▶ $q_2(s) = \mathcal{O}_{s \rightarrow 0^+}(q_1(s))$: Given $q_1, q_2 \in \mathcal{PD}$, we say that q_1 *has greater growth* than q_2 around zero if $\exists k \geq 0$ such that $\limsup_{s \rightarrow 0^+} q_2(s)/q_1(s) \leq k$.

Cascades of Strong iISS Systems



Theorem: Strong iISS is preserved under cascade [Chaillet et. al., Automatica, 2014]

If the systems $\dot{x}_1 = f_1(x_1, u_1)$ and $\dot{x}_2 = f_2(x_2, u_2)$ are **Strongly iISS**, then the cascade (2) is **Strongly iISS**.

Corollary: iISS + Strong iISS \Rightarrow iISS [Chaillet et. al., Automatica, 2014]

If $\dot{x}_1 = f_1(x_1, u_1)$ is **Strongly iISS** and $\dot{x}_2 = f_2(x_2, u_2)$ is **iISS**, then (2) is **iISS**.

Corollary: GAS + Strong iISS \Rightarrow GAS [Chaillet et. al., Automatica, 2014]

If $\dot{x}_1 = f_1(x_1, u_1)$ is **Strongly iISS** and $\dot{x}_2 = f_2(x_2)$ is **GAS**, then

$$\begin{aligned}\dot{x}_1 &= f_1(x_1, x_2) \\ \dot{x}_2 &= f_2(x_2)\end{aligned} \quad \text{is GAS.}$$

ISS/iISS Systems: Summary

	ISS	Strong iISS	iISS
0-GAS	✓	✓	✓
Forward completeness $\forall u \in \mathcal{U}$	✓	✓	✓
Bounded input-Bounded state	✓	For $\ u\ < R$	☹
Converging input-Converging state	✓	✓	☹
Preservation under cascade	✓	✓	Growth rate
Lyapunov characterization	$\alpha \in \mathcal{K}_\infty$	Open question	$\alpha \in \mathcal{PD}$